




Abstract only
PubMed



PubMed

Nucleotide

Protein

Genome

Structure

PopSet

Taxonomy

OMIM

Bc

Search

PubMed



for



Limits

Preview/Index

History

Clipboard

Details

About Entrez



Abstract



Sort



Save

Text

Clip/Add

Order

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

PubMed Services

Journal Browser

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Mol Cell Biol 1994 Sep;14(9):5671-81 Related Articles, **NEW Books**, LinkOut

The lung-specific surfactant protein B gene promoter is a target for thyroid transcription factor 1 and hepatocyte nuclear factor 3, indicating common factors for organ-specific gene expression along the foregut axis.

Bohinski RJ, Di Lauro R, Whitsett JA.

Division of Pulmonary Biology, Children's Hospital Medical Center, Cincinnati, Ohio 45229-2899.

We used the lung epithelial cell-specific surfactant protein B (SPB) gene promoter as a model with which to investigate mechanisms involved in transcriptional control of lung-specific genes. In a previous study, we showed that the SPB promoter specifically activated expression of a linked reporter gene in the continuous H441 lung cell line and that H441 nuclear proteins specifically protected a region of this promoter from bp -111 to -73. In this study, we further show that this region is a complex binding site for thyroid transcription factor 1 (TTF-1) and hepatocyte nuclear factor 3 (HNF-3). Whereas TTF-1 bound two highly degenerate and closely spaced sites, HNF-3 proteins bound a TGT3 motif (TGTTTGT) that is also found in several liver-specific gene regulatory regions, where it appears to be a weak affinity site for HNF-3. Point mutations of these binding sites eliminated factor binding and resulted in significant decreases in transfected SPB promoter activity. In addition, we developed a cotransfection assay and showed that a family of lung-specific gene promoters that included the SPB, SPC, SPA, and Clara cell secretory protein (CCSP) gene promoters were specifically activated by cotransfected TTF-1. We conclude that TTF-1 and HNF-3 are major activators of lung-specific genes and propose that these factors are involved in a general mechanism of lung-specific gene transcription. Importantly, these data also show that common factors are involved in organ-specific gene expression along the mammalian foregut axis.

PMID: 8065304 [PubMed - indexed for MEDLINE]